arXiv:1510.01849v1 [math.AG] 7 Oct 2015

Isomorphism classes of Doche-Icart-Kohel Curves over finite fields

Reza Rezaeian Farashahi
Department of Mathematical Sciences
Isfahan University of Technology
P.O. Box 85145Isfahan, Iran
and
School of Mathematics,
Institute for Research in Fundamental Sciences (IPM),
P.O. Box 19395-5746, Tehran, Iran
farashahi@cc.iut.ac.ir

Mehran Hosseini
Department of Mathematical Sciences
Isfahan University of Technology
mehran.hosseini@math.iut.ac.ir

Abstract

We give explicit formulas for the number of distinct elliptic curves over a finite field, up to isomorphism, in two families of curves introduced by C. Doche, T. Icart and D. R. Kohel.

Keywords: Elliptic curve, j-invariant, isomorphism, cryptography

2000 Mathematics Subject Classification: 11G05,11T06, 14H52

1 Introduction

An elliptic curve is a smooth projective genus 1 curve, with a given rational point. Traditionally, an elliptic curve E over a filed $\mathbb{F}$ is represented by the Weierstrass equation

$$E: \quad Y^2 + a_1XY + a_3Y = X^3 + a_2X^2 + a_4X + a_6, \tag{1}$$

where the coefficients $a_1, a_2, a_3, a_4, a_6 \in \mathbb{F}$. Elliptic curves can be represented by several other models (see [1, Chapter 13] or [8, Chapter 2]). In the past few years, these alternative models have been revisited duo to cryptoraphic applications. Moreover, some new families of elliptic curves have been proposed following the cryptographic interests. In the cryptographic settings the curves are usually considered over finite fields $\mathbb{F}_q$ of q element.

In this work, we first consider the family of elliptic curves introduced by C. Doche, T. Icart and D. R. Kohel [2] over finite fields $\mathbb{F}_q$ of characteristic $p \geq 5$

$$\mathbf{E}_{\mathrm{T},u}: \quad Y^2 = X^3 + 3u(X+1)^2, \tag{2}$$

where $u \neq 0, 9/4$. Doche et. al. have build this family of elliptic curves for which the isogeny of multiplication by three splits into 2 isogenies of degree 3. In this way, they proposed more efficient tripling formulas leading to a fast scalar multiplication algorithm. Notice, an elliptic curve defined over $\mathbb{F}_q$ with a rational 3-torsion subgroup can be expressed in the latter form (up to twists). They also proposed another family

$$\mathbf{E}_{\mathrm{D},u}: \quad Y^2 = X^3 + uX^2 + 16uX, \tag{3}$$

where $u \neq 0, 64$ and $p \geq 3$. This family represents elliptic curves defined over $\mathbb{F}_q$ with a rational 2-torsion subgroup (up to twists). Accordingly a natural question arises about the number of distinct (up to isomorphism) elliptic curves over $\mathbb{F}_q$ in above families (2) and (3).

Farashahi and Shparlinski [5], using the notion of the *j-invariant of an elliptic curve*, see [1, 7, 8], gave *exact* formulas for the number of distinct elliptic curves $\mathbf{E}_{\mathrm{T},u}$ over $\mathbb{F}_q$ (up to isomorphism over the algebraic closure of $\mathbb{F}_q$) when $u \in \mathbb{F}_q \setminus \{0, 9/4\}$. In addition, the exact formulas for the numbers of distinct elliptic curves over a finite field $\mathbb{F}_q$ in several families are provided in [3, 4]. We notice, the formulas given in [5, 3, 4] are exactly given in terms of q. Accordingly, the open question of [5] is whether there is explicit formula for the number of distinct elliptic curves (up to isomorphism over $\mathbb{F}_q$) in the family (2). We answer this question with explicit estimates for the number of $\mathbb{F}_q$-isomorphism classes in the families of Doche, Icart and Kohel. The interesting point is the numbers of $\mathbb{F}_q$-isomorphism classes of the families (2) and (3) depends not only on q but also on the number of $\mathbb{F}_q$ rational points of some corresponding curve. In fact, we have interesting examples of families of elliptic curves where the number of $\mathbb{F}_q$-isomorphism classes of the family is given approximately by Hasse-Weil bound. Our counting proof techniques are slightly different in both families (2) and (3).

Throughout the paper, for a field $\mathbb{F}$, we denote its algebraic closure by $\overline{\mathbb{F}}$ and its multiplicative subgroup by $\mathbb{F}^*$. The letter p always denotes a prime number and the letter q always denotes a prime power of p. As usual, $\mathbb{F}_q$ is a finite field of size q. Let χ_1 denote the principal character in $\mathbb{F}_q$ where $p \geq 5$. Let χ_2 denote the quadratic character in $\mathbb{F}_q$, where $p \geq 3$. Then, for any q where $p \geq 3$, we have $u = w^2$ for some $w \in \mathbb{F}_q^*$ if and only if $\chi_2(u) = 1$. And, let χ_3 denote a primitive cubic character in $\mathbb{F}_q$ if $q \equiv 1 \pmod 3$ and $\chi_3 = \chi_1$ if $q \equiv 2 \pmod 3$. In particular, for any field $\mathbb{F}_q$ of characteristic $p \geq 5$, $u = w^3$ for some $w \in \mathbb{F}_q^*$ if and only if $\chi_3(u) = 1$. We recall that $U = O(V)$ is equivalent to the

inequality $|U| \leq cV$ with some constant $c > 0$. The cardinality of a finite set $\mathcal{S}$ is denoted by $\#\mathcal{S}$.

2 Isomorphisms of elliptic curves

Here, we briefly recall some words on isomorphisms between elliptic curves, see [1, 7, 8] for a general background on isomorphisms and elliptic curves. Two elliptic curves given by Weierstrass equations (1) are isomorphic over a field $\mathbb{F}$ if and only if there is a change of variables between them of the form:

$$(x, y) \to (\alpha^2 x + r, \alpha^3 y + \alpha^2 s x + t),$$

where $\alpha \neq 0$, and $\alpha, r, s, t \in \mathbb{F}$. We use $E_1 \cong_{\mathbb{F}} E_2$ to denote the elliptic curves E_1 and E_2 are $\mathbb{F}$-*isomorphic*. If $\alpha, r, s, t \in \overline{\mathbb{F}}$, the two elliptic curves are called *isomorphic* over $\overline{\mathbb{F}}$ or *twists* of each other.

The elliptic curve E over $\mathbb{F}$ given by the Weierstrass equation (1) has the non-zero discriminant

$$\Delta_E = -b_2^2 b_8 - 8b_4^3 - 27b_6^2 + 9b_2 b_4 b_6,$$

$$b_2 = a_1^2 + 4a_2,\ b_4 = a_1 a_3 + 2a_4,\ b_6 = a_3^2 + 4a_6,\ b_8 = a_1^2 a_6 - a_1 a_3 a_4 + 4a_2 a_6 + a_2 a_3^2 - a_4^2.$$

And, the j-invariant of E is explicitly defined as

$$j(E) = (b_2^2 - 24b_4)^3 / \Delta_E.$$

It is known that two elliptic curves E_1, E_2 over a field $\mathbb{F}$ are isomorphic over $\overline{\mathbb{F}}$ if and only if $j(E_1) = j(E_2)$, see [7, Proposition III.1.4(b)]. However, two elliptic curves with the same j-invariant need not be isomorphic over $\mathbb{F}$.

Any elliptic curve over a finite field $\mathbb{F}_q$ of characteristic $p \geq 5$ can be represented by the short Weierstrass equation

$$E_{a,b}: \quad Y^2 = X^3 + aX + b,$$

where the coefficients $a, b \in \mathbb{F}_q$, and $\Delta_{E_{a,b}} = -16(4a^3 + 27b^2) \neq 0$. The the j-invariant of $E_{a,b}$ is given by

$$j(E_{a,b}) = 1728 \frac{4a^3}{4a^3 + 27b^2}.$$

Two elliptic curves E_{a_1,b_1}, E_{a_2,b_2} represented in the short Weierstrass form are isomorphic over $\mathbb{F}_q$ if and only if there exists a nonzero element $\alpha \in \mathbb{F}_q$ with the change of variables between them of the form

$$(x, y) \to (\alpha^2 x, \alpha^3 y).$$

In other words, $E_{a_1,b_1} \cong_{\mathbb{F}_q} E_{a_2,b_2}$ if and only if there exists a nonzero element α in $\mathbb{F}_q$ such that $a_1 = \alpha^4 a_2$ and $b_1 = \alpha^6 b_2$. The short Weierstrass model of the elliptic curve $\mathbf{E}_{\mathrm{T},u}$ given by the equation (2) is represented by

$$\mathbf{E}_{\mathrm{W},u}: \quad Y^2 = X^3 + a_u X + b_u, \tag{4}$$

where $a_u = -2^4 3^5 u(u-2)$, $b_u = 2^6 3^6 u(2u^2 - 6u + 3)$. The j-invariant of $\mathbf{E}_{\mathrm{T},u}$ is obtained as

$$j(\mathbf{E}_{\mathrm{T},u}) = \frac{6912u(u-2)^3}{4u-9}.$$

Similarly, the j-invariant of $\mathbf{E}_{\mathrm{D},u}$ given by the equation (3) is obtained as

$$j(\mathbf{E}_{\mathrm{D},u}) = \frac{(u-48)^3}{u-64}.$$

We use $J_{\mathrm{T}}(q)$ and $J_{\mathrm{D}}(q)$ to denote the numbers of distinct j-invariants of the elliptic curves defined over $\mathbb{F}_q$ in the families (2) and (3) respectively. We have, [5, Theorem 9],

$$J_{\mathrm{T}}(q) = \lfloor (3q+1)/4 \rfloor \text{ if } q \equiv 1 \pmod 3 \quad \text{and} \quad J_{\mathrm{T}}(q) = (q-1)/2 \text{ if } q \equiv 2 \pmod 3.$$

In Theorem (5) we provide the number $J_{\mathrm{D}}(q)$ for any finite field $\mathbb{F}_q$ of characteristic $p \geq 3$.

An $\overline{\mathbb{F}}_q$-isomorphism class of elliptic curves in a given family may be an $\mathbb{F}_q$-isomorphism class or divide into two or more $\mathbb{F}_q$-isomorphism classes. The former occurs when all elliptic curve in the $\overline{\mathbb{F}}_q$-isomorphism class are $\mathbb{F}_q$-isomorphic to each others, and the latter occurs if at least two elliptic curves in the class are not $\mathbb{F}_q$-isomorphic, i.e., this happens if there are non trivial quadratic twists in the $\overline{\mathbb{F}}_q$-isomorphism class. Therefore, in a given family the number of $\mathbb{F}_q$-isomorphism classes may not be so related to the number of $\overline{\mathbb{F}}_q$-isomorphism classes. In this paper, we use $I_{\mathrm{T}}(q)$ and $I_{\mathrm{T}}(q)$ to denote the numbers of $\mathbb{F}_q$-isomorphism classes of the families (2) and (3) respectively. And, in the next sections we give explicit formulas for these numbers.

3 Tripling Doche-Icart-Kohel curves

3.1 Preliminaries

We consider the curves $\mathbf{E}_{\mathrm{T},u}$ given by (2) over a finite field $\mathbb{F}_q$ of characteristic $p \geq 5$. Notice, $u \notin \{0, 9/4\}$, since the curve $\mathbf{E}_{\mathrm{T},u}$ is nonsingular. Then, the j-invariant $j(\mathbf{E}_{\mathrm{T},u}) = F(u)$ where

$$F(U) = \frac{2^8 3^3 U(U-2)^3}{4U-9}. \tag{5}$$

We recall that two elliptic the curves $\mathbf{E}_{\mathrm{T},u}$ and $\mathbf{E}_{\mathrm{T},v}$ are isomorphic over $\overline{\mathbb{F}}_q$ if and only if they have the same j-invariants. So, the number of distinct elliptic curves $\mathbf{E}_{\mathrm{T},u}$ up to isomorphism over $\overline{\mathbb{F}}_q$ equals the number of distinct values $F(u)$, for all $u \in \mathbb{F}_q \setminus \{0, 9/4\}$. To compute this number, we consider the bivariate rational function $F(U) - F(V) = g(U,V)/h(U,V)$ with two relatively prime polynomials g and h. We see that

$$\begin{aligned} g(U,V) =& 4U^3V - 9U^3 + 4U^2V^2 - 33U^2V + 54U^2 + \\ & 4UV^3 - 33UV^2 + 102UV - 108U - 9V^3 + 54V^2 - 108V + 72. \end{aligned}$$

For a fixed value $u \in \mathbb{F}_q \setminus \{0, 9/4\}$, let $g_u(V) = g(u, V)$ and let

$$\Delta_u = -12(u(u-2)(4u-9)(2u^2-6u+3))^2$$

be the discriminant of g_u. Furthermore, we let

$$\mathcal{Z}_u = \{v \ : \ v \in \mathbb{F}_q \setminus \{0, 9/4\}, \ v \neq u, \ g_u(v) = 0\}.$$

Moreover, for $u \in \mathbb{F}_q \setminus \{0, 9/4\}$, we let

$$\mathcal{J}_{\mathrm{T},u} = \left\{v \ : \ v \in \mathbb{F}_q \setminus \{0, 9/4\}, \quad \mathbf{E}_{\mathrm{T},u} \cong_{\overline{\mathbb{F}}_q} \mathbf{E}_{\mathrm{T},v}\right\}.$$

Clearly, we have $\mathcal{J}_{\mathrm{T},u} = \mathcal{Z}_u \cup \{u\}$. We let

$$\psi(U) = \frac{2(4U-9)}{U}.$$

Recall from [5, Lemma 6], that for $u \in \mathbb{F}_q \setminus \{0, 9/4\}$ with $\Delta_u \neq 0$ we obtain

$$\mathcal{Z}_u = \left\{v \ : \ v \in \mathbb{F}_q \setminus \{0, 9/4\}, v = -\frac{1}{3}\left(u - 6 - wu + \frac{2(u-3)}{w}\right), \ w \in \mathbb{F}_q, \ w^3 = \psi(u)\right\}. \tag{6}$$

For the particular case of $\Delta_u = 0$, we have $\mathcal{J}_{\mathrm{T},u} = \{u\}$ if $u = 2$ and $\mathcal{J}_{\mathrm{T},u} = \{u, -u+3\}$ if $2u^2 - 6u + 3 = 0$. The former case occurs for the elliptic curve $\mathbf{E}_{\mathrm{T},u}$ with $j(\mathbf{E}_{\mathrm{T},u}) = 0$ and the latter case corresponds with elliptic curves $\mathbf{E}_{\mathrm{T},u}$ with $j(\mathbf{E}_{\mathrm{T},u}) = 1728$.

We partition the set $\mathbb{F}_q \setminus \{0, 9/4\}$ into the following sets:

$$\mathcal{A}_1 \cup \mathcal{A}_2 \cup \mathcal{A}_3 \cup \mathcal{B}_1 \cup \mathcal{B}_2, \tag{7}$$

where

$$\begin{aligned}
\mathcal{A}_1 &= \{u \in \mathbb{F}_q \setminus \{0, 9/4\} \ : \ \Delta_u \neq 0, \ \chi_3(\psi(u)) \neq 1 \text{ and } q \equiv 1 \pmod 3\}, \\
\mathcal{A}_2 &= \{u \in \mathbb{F}_q \setminus \{0, 9/4\} \ : \ \Delta_u \neq 0 \text{ and } q \equiv 2 \pmod 3\}, \\
\mathcal{A}_3 &= \{u \in \mathbb{F}_q \setminus \{0, 9/4\} \ : \ \Delta_u \neq 0, \ \chi_3(\psi(u)) = 1 \text{ and } q \equiv 1 \pmod 3\}, \\
\mathcal{B}_1 &= \{u = 2\}, \\
\mathcal{B}_2 &= \left\{u \in \mathbb{F}_q \setminus \{0, 9/4\} \ : \ 2u^2 - 6u + 3 = 0\right\}.
\end{aligned}$$

The cardinalities of above sets are provided in Table 1 [5, Table 7], where we let $q \equiv r \pmod{12}$.

Furthermore, from [5, Lemmas 7,8], for all $u \in \mathbb{F}_q \setminus \{0, 9/4\}$ we have

$$\#\mathcal{J}_{\mathrm{T},u} = \begin{cases} 1, & \text{if } u \in \mathcal{A}_1 \cup \mathcal{B}_1, \\ 2, & \text{if } u \in \mathcal{A}_2 \cup \mathcal{B}_2, \\ 4, & \text{if } u \in \mathcal{A}_3. \end{cases} \tag{8}$$

r	$\#\mathcal{A}_1$	$\#\mathcal{A}_2$	$\#\mathcal{A}_3$	$\#\mathcal{B}_1$	$\#\mathcal{B}_2$
1	$\frac{2(q-1)}{3}$	0	$\frac{q-1}{3}-4$	1	2
5	0	$q-3$	0	1	0
7	$\frac{2(q-1)}{3}$	0	$\frac{q-1}{3}-2$	1	0
11	0	$q-5$	0	1	2

Table 1: Cardinalities of the sets $\mathcal{A}_1, \mathcal{A}_2, \mathcal{A}_3, \mathcal{B}_1, \mathcal{B}_2$.

Next, for the number $J_{\mathrm{T}}(q)$ of $\overline{\mathbb{F}}_q$-isomorphism class of elliptic curves in the family (2), that is the cardinality of the image of the polynomial F given by (5), we have

$$J_{\mathrm{T}}(q) = \#\mathcal{A}_1 + \#\mathcal{B}_1 + \frac{\#\mathcal{A}_2 + \#\mathcal{B}_2}{2} + \frac{\#\mathcal{A}_3}{4}.$$

Next, using Table 1, a precise formula for the number of distinct curves of the family (2) is obtained (see [5, Theorem 9]).

Theorem 1. *For any finite field $\mathbb{F}_q$ of characteristic $p \geq 5$, for the number $J_{\mathrm{T}}(q)$ of distinct values of the j-invariant of the family (2), we have*

$$J_{\mathrm{T}}(q) = \begin{cases} \left\lfloor \dfrac{3q+1}{4} \right\rfloor & \text{if } q \equiv 1 \pmod 3, \\ \dfrac{q-1}{2} & \text{if } q \equiv 2 \pmod 3. \end{cases}$$

Notice, that elliptic curves $\mathbf{E}_{\mathrm{T},u}$ and $\mathbf{E}_{\mathrm{T},v}$, where $v \in \mathcal{J}_{\mathrm{T},u}$, may not be isomorphic over $\mathbb{F}_q$. So, for a fixed value $u \in \mathbb{F}_q \setminus \{0, 9/4\}$, we let

$$\mathcal{I}_{\mathrm{T},u} = \left\{ v \ : \ v \in \mathbb{F}_q \setminus \{0, 9/4\}, \ \mathbf{E}_{\mathrm{T},u} \cong_{\mathbb{F}_q} \mathbf{E}_{\mathrm{T},v} \right\}.$$

Clearly, for all $u \in \mathbb{F}_q \setminus \{0, 9/4\}$, we have $\mathcal{I}_{\mathrm{T},u} \subseteq \mathcal{J}_{\mathrm{T},u}$. To count the number of distinct $\mathbb{F}_q$-isomorphism classes of elliptic curves $\mathbf{E}_{\mathrm{T},u}$, for all $u \in \mathbb{F}_q \setminus \{0, 9/4\}$, we need to find the number of distinct sets $\mathcal{I}_{\mathrm{T},u}$. Therefore, first we shall obtain the cardinalities of the sets $\mathcal{I}_{\mathrm{T},u}$, for all $u \in \mathbb{F}_q \setminus \{0, 9/4\}$.

We recall that the short Weierstrass model of the elliptic curve $\mathbf{E}_{\mathrm{T},u}$ is given by (4) as

$$\mathbf{E}_{\mathrm{W},u} : \quad Y^2 = X^3 + a_u X + b_u,$$

where $a_u = -2^4 3^5 u(u-2)$, $b_u = 2^6 3^6 u(2u^2 - 6u + 3)$.

Let $u \in \mathbb{F}_q \setminus \{0, 9/4\}$ and let $v \in \mathcal{I}_{\mathrm{T},u}$, where $v \neq u$. Therefore, $v \in \mathcal{J}_{\mathrm{T},u}$. So, from (6) we have, $v = -\left(u - 6 - wu + 2(u-3)/w\right)/3$, where $w^3 = \psi(u) = 2(4u-9)/u$. Since the elliptic curves $\mathbf{E}_{\mathrm{T},u}$ and $\mathbf{E}_{\mathrm{T},v}$ are isomorphic over $\mathbb{F}_q$, their Weierstrass models $\mathbf{E}_{\mathrm{W},u}$ and $\mathbf{E}_{\mathrm{W},v}$ are $\mathbb{F}_q$-isomorphic. We have

$$a_v = \left(\frac{(w+1)(w-2)}{3w}\right)^2 a_u, \qquad b_v = \left(\frac{(w+1)(w-2)}{3w}\right)^3 b_u.$$

This means, if $\mathbf{E}_{\mathrm{W},u} \cong_{\mathbb{F}_q} \mathbf{E}_{\mathrm{W},v}$ then $\frac{(w+1)(w-2)}{3w}$ is a quadratic residue in $\mathbb{F}_q$.

Therefore, if $\mathbf{E}_{\mathrm{T},u} \cong_{\mathbb{F}_q} \mathbf{E}_{\mathrm{T},v}$ then w is the nonzero x-coordinate of an $\mathbb{F}_q$-rational point on the elliptic curve

$$E : Y^2 = 3X(X+1)(X-2).$$

On the other hand, if w is the nonzero x-coordinate of an $\mathbb{F}_q$-rational point on the elliptic curve E and if we let u, v be distinct elements of $\mathbb{F}_q \setminus \{0, 9/4\}$ such that

$$\psi(u) = w^3, \quad v = -\frac{1}{3}(u - 6 - wu + \frac{2(u-3)}{w}),$$

then $\mathbf{E}_{\mathrm{T},u} \cong_{\mathbb{F}_q} \mathbf{E}_{\mathrm{T},v}$. In other words, we have the following Lemma 2.

Lemma 2. *For all distinct elements $u, v \in \mathbb{F}_q \setminus \{0, 9/4\}$, we have $\mathbf{E}_{\mathrm{T},u} \cong_{\mathbb{F}_q} \mathbf{E}_{\mathrm{T},v}$ if and only if we have*

$$u = \frac{-18}{w^3 - 8} \qquad \text{and} \qquad v = \frac{2(w+1)^3}{w(w^2+2w+4)},$$

for some $w \in \mathbb{F}_q$ with $\chi_2\left(3w(w+1)(w-2)\right) = 1$.

3.2 The number of curves in the family (2)

In this section, we look into the number of $\mathbb{F}_q$-isomorphism classes of tripling Doche-Icart-Kohel curves. The estimate for the number $I_{\mathrm{T}}(q)$ of $\mathbb{F}_q$-isomorphism classes of this family is given by

$$I_{\mathrm{T}}(q) = \begin{cases} (79/96)q + O(\sqrt{q}) & \text{if } q \equiv 1 \pmod 3, \\ (3/4)q + O(\sqrt{q}) & \text{if } q \equiv 2 \pmod 3. \end{cases}$$

Furthermore, in the next theorem we provide explicit formulas for the number of $\mathbb{F}_q$-isomorphism classes of Doche-Icart-Kohel curves.

Theorem 3. *For any finite field $\mathbb{F}_q$ of characteristic $p \geq 5$, for the number $I_{\mathrm{T}}(q)$ of $\mathbb{F}_q$-isomorphism classes of the family (2), we have*

$$I_{\mathrm{T}}(q) = \begin{cases} (5q-2)/6 - (N_1 - 25)/96 & \text{if } q \equiv 1 \pmod{24}, \\ (5q+1)/6 - (N_1 - 25)/96 & \text{if } q \equiv 13 \pmod{24}, \\ 5(q-1)/6 - (N_1 - 25)/96 & \text{if } q \equiv 7 \pmod{12}, \\ q - 1 - N_2/4 & \text{if } q \equiv 2 \pmod 3, \end{cases}$$

where N_1 is the number of $\mathbb{F}_q$-rational affine points of the curve $\mathcal{C}$ given by

$$\begin{cases} Y^2 = 3X(X+1)(X-2), \\ Z^2 = 3\zeta X(\zeta X+1)(\zeta X-2), \\ W^2 = 3\zeta^2 X(\zeta^2 X+1)(\zeta^2 X-2), \end{cases}$$

where ζ is a nontrivial cubic root of the unity in $\mathbb{F}_q$ where $q \equiv 1 \pmod 3$, and N_2 is the number of $\mathbb{F}_q$-rational points of the Legendre curve

$$\mathcal{L} : Y^2 = X(X-1)(X-1/3).$$

Proof. Let $\mathcal{J}_{\mathrm{T}}$ be the set of j-invariants of the elliptic curve $\mathbf{E}_{\mathrm{T},u}$ for all $u \in \mathbb{F}_q \setminus \{0, 9/4\}$. For $a \in \mathbb{F}_q$, let $i_{\mathrm{T}}(a)$ be the set of distinct $\mathbb{F}_q$-isomorphism classes of elliptic curves $\mathbf{E}_{\mathrm{T},u}$ with $j(\mathbf{E}_{\mathrm{T},u}) = a$. Clearly, $\#i_{\mathrm{T}}(a) = 0$, if $a \notin \mathcal{J}_{\mathrm{T}}$. Therefore, we have

$$I_{\mathrm{T}}(q) = \sum_{a \in \mathbb{F}_q} \#i_{\mathrm{T}}(a) = \sum_{a \in \mathcal{J}_{\mathrm{T}}} \#i_{\mathrm{T}}(a) \ .$$

Notice that, for $a \in \mathcal{J}_{\mathrm{T}}$, the cardinality of the set $i_{\mathrm{T}}(a)$ equals 1 or 2. Let $\mathcal{T}$ be the set of all $a \in \mathcal{J}_{\mathrm{T}}$ where all elliptic curves $\mathbf{E}_{\mathrm{T},u}$ with j-invariant a are $\mathbb{F}_q$-isomorphic to each others. In other words, we have

$$\mathcal{T} = \{a \ : \ a \in \mathcal{J}_{\mathrm{T}} \ , \ \#i_{\mathrm{T}}(a) = 1\} \, .$$

Then, $\#i_{\mathrm{T}}(a) = 2$ if $a \in \mathcal{J}_{\mathrm{T}} \setminus \mathcal{T}$. Thus,

$$\begin{aligned} I_{\mathrm{T}}(q) = \sum_{a \in \mathcal{J}_{\mathrm{T}}} \#i_{\mathrm{T}}(a) &= \sum_{a \in \mathcal{T}} \#i_{\mathrm{T}}(a) + \sum_{a \in \mathcal{J}_{\mathrm{T}} \setminus \mathcal{T}} \#i_{\mathrm{T}}(a) \\ &= \sum_{a \in \mathcal{T}} 1 + \sum_{a \in \mathcal{J}_{\mathrm{T}} \setminus \mathcal{T}} 2 = \#\mathcal{T} + 2(\#\mathcal{J}_{\mathrm{T}} - \#\mathcal{T}) \\ &= 2J_{\mathrm{T}}(q) - \#\mathcal{T} \ . \end{aligned}$$

Next, we compute the cardinality of the set $\mathcal{T}$. We recall the partition of the set $\mathbb{F}_q \setminus \{0, 9/4\}$ into

$$\mathcal{A}_1 \cup \mathcal{A}_2 \cup \mathcal{A}_3 \cup \mathcal{B}_1 \cup \mathcal{B}_2,$$

given by (7). For $i = 1, 2, 3$, let

$$\widetilde{\mathcal{A}}_i = \{u \ : \ u \in \mathcal{A}_i, \ j(\mathbf{E}_{\mathrm{T},u}) \in \mathcal{T}\} \, ,$$

and for $i = 1, 2$ we let

$$\widetilde{\mathcal{B}}_i = \{u \ : \ u \in \mathcal{B}_i, \ j(\mathbf{E}_{\mathrm{T},u}) \in \mathcal{T}\} \, .$$

Then, from (8), we have

$$\#\mathcal{T} = \#\widetilde{\mathcal{A}}_1 + \#\widetilde{\mathcal{B}}_1 + \frac{\#\widetilde{\mathcal{A}}_2 + \#\widetilde{\mathcal{B}}_2}{2} + \frac{\#\widetilde{\mathcal{A}}_3}{4}. \tag{9}$$

If $u = 2 \in \mathcal{B}_1$ then $j(\mathbf{E}_{\mathrm{T},u}) = 0$ and $\mathcal{J}_{\mathrm{T},u} = \{u\}$. Thus, $\#i_{\mathrm{T}}(0) = 1$ and $\#\widetilde{\mathcal{B}}_1 = 1$. If $u \in \mathcal{B}_2$ i.e., $2u^2 - 6u + 3 = 0$, then $j(\mathbf{E}_{\mathrm{T},u}) = 1728$ and $\mathcal{J}_{\mathrm{T},u} = \{u, -u + 3\}$. The short Weierstrass model of $\mathbf{E}_{\mathrm{T},u}$ is given by (4) as $\mathbf{E}_{\mathrm{W},u} : \ Y^2 = X^3 + a_u X$, where $a_u = -2^4 3^5 u(u-2)$. We see that $a_u = -a_{-u+3}$. So, the elliptic curves $\mathbf{E}_{\mathrm{T},u}$ and $\mathbf{E}_{\mathrm{T},-u+3}$ are $\mathbb{F}_q$-isomorphic if and only if $-1 = w^4$ for some $w \in \mathbb{F}_q^*$. The latter is equivalent to have $q \equiv 1 \pmod 8$. Also, we recall from Table 1, that

$$\#\mathcal{B}_2 = \begin{cases} 2, & \text{if } q \equiv \pm 1 \pmod{12}, \\ 0, & \text{if } q \not\equiv \pm 1 \pmod{12}. \end{cases}$$

So, we have

$$\#i_{\mathrm{T}}(1728) = \begin{cases} 2, & \text{if } q \equiv 11, 13, 23 \pmod{24}, \\ 1, & \text{if } q \equiv 1 \pmod{24}, \\ 0, & \text{if} q \not\equiv \pm 1 \pmod{12}, \end{cases}$$

and

$$\#\widetilde{\mathcal{B}}_2 = \begin{cases} 2, & \text{if } q \equiv 1 \pmod{24}, \\ 0, & \text{if } q \not\equiv 1 \pmod{24}. \end{cases}$$

If $u \in \mathcal{A}_1$, by (8) we have $\#\mathcal{J}_{\mathrm{T},u} = 1$. So, $\#i_{\mathrm{T}}(a) = 1$, where $a = j(\mathbf{E}_{\mathrm{T},u})$. Therefore, for all $u \in \mathcal{A}_1$ we have $j(\mathbf{E}_{\mathrm{T},u}) \in \mathcal{T}$. Thus

$$\widetilde{\mathcal{A}}_1 = \mathcal{A}_1.$$

If $u \in \mathcal{A}_2$, then $u \in \mathbb{F}_q \backslash \{0, 2, 9/4\}$ where $q \equiv 2 \pmod 3$, and $2u^2 - 6u + 3 \neq 0$. By (8) we have $\#\mathcal{J}_{\mathrm{T},u} = 2$. More precisely, we have $\mathcal{J}_{\mathrm{T},u} = \{u, v\}$, where $v = -\frac{1}{3}(u - 6 - wu + \frac{2(u-3)}{w})$ and $w^3 = \psi(u) = 2(4u - 9)/u$ where $w \in \mathbb{F}_q$. Notice that w is uniquely determined by u, since $q \equiv 2 \pmod 3$. Furthermore, from Lemma 2, $\mathbf{E}_{\mathrm{T},u} \cong_{\mathbb{F}_q} \mathbf{E}_{\mathrm{T},v}$ if and only if we have $\chi_2\left(3w(w+1)(w-2)\right) = 1$.So, $j(\mathbf{E}_{\mathrm{T},u}) \in \mathcal{T}$ if and only if $\chi_2\left(3w(w+1)(w-2)\right) = 1$. Therefore,

$$\widetilde{\mathcal{A}}_2 = \left\{ u \ : \ u \in \mathcal{A}_2, \ u = \frac{-18}{w^3 - 8}, \ \chi_2\left(3w(w+1)(w-2)\right) = 1 \right\}.$$

Let L be the elliptic curve over $\mathbb{F}_q$ given by $L : Y^2 = 3X(X+1)(X-2)$. Let $P = (x, y)$be a point of $L(\mathbb{F}_q)$ with $y \neq 0$. Let $u = \frac{-18}{x^3-8}$. Then, $u \neq 0, 2, 9/4$. Furthermore, $2u^2-6u+3 \neq 0$. Since if $2u^2 - 6u + 3 = 0$, then 3 is a quadratic residue in $\mathbb{F}_q$ where $q \equiv 11 \pmod{12}$. And, we have $x^2 + 2x - 2 = 0$ and $3x(x+1)(x-2) = -9x^2$, which contradicts $\chi_2(-1) = -1$. Therefore, $u \in \mathcal{A}_2$ and $u \in \widetilde{\mathcal{A}}_2$. In other word, for $q \equiv 2 \pmod 3$,

$$\widetilde{\mathcal{A}}_2 = \left\{ u \ : \ u = \frac{-18}{x^3 - 8}, \ (x, y) \in L(\mathbb{F}_q), y \neq 0 \right\}.$$

The latter means, $\#\widetilde{\mathcal{A}}_2 = (\#L(\mathbb{F}_q) - 4)/2$ if $q \equiv 2 \pmod 3$. In addition,the elliptic curve L is $\mathbb{F}_q$-isomorphic to the Legendre curve $\mathcal{L}$ and we have

$$\#\widetilde{\mathcal{A}}_2 = \begin{cases} 0, & \text{if } q \equiv 1 \pmod 3 \\ (\#\mathcal{L}(\mathbb{F}_q) - 4)/2, & \text{if } q \equiv 2 \pmod 3 \end{cases}$$

If $u \in \mathcal{A}_3$, then $u \in \mathbb{F}_q \backslash \{0, 2, 9/4\}$ where $q \equiv 1 \pmod 3$, and $2u^2 - 6u + 3 \neq 0$, $\chi_3(\psi(u)) = 1$. From (8) we have $\#\mathcal{J}_{\mathrm{T},u} = 4$. So, we write $\mathcal{J}_{\mathrm{T},u} = \{u, v_1, v_2, v_3\}$, where for $i = 1, 2, 3$, $v_i = -\frac{1}{3}(u - 6 - w_i u + \frac{2(u-3)}{w_i})$ and $w_i^3 = \psi(u) = 2(4u - 9)/u$. Here, w_1, w_2, w_3 are cubic roots of $\psi(u)$ in $\mathbb{F}_q$. By Lemma 2, $\mathbf{E}_{\mathrm{T},u} \cong_{\mathbb{F}_q} \mathbf{E}_{\mathrm{T},v_i}$ if and only if we have $\chi_2\left(3w_i(w_i+1)(w_i-2)\right) = 1$. So, $j(\mathbf{E}_{\mathrm{T},u}) \in \mathcal{T}$ if and only if $\chi_2\left(3w_i(w_i+1)(w_i-2)\right) = 1$ for $i = 1, 2, 3$. For simplicity, we let $w = w_1$. And, we have $w_2 = \zeta w$, $w_3 = \zeta^2 w$. Therefore,

$$\widetilde{\mathcal{A}}_3 = \left\{ u \ : \ u \in \mathcal{A}_3, \ u = \frac{-18}{w^3 - 8}, \ \chi_2\left(3\zeta^i w(\zeta^i w + 1)(\zeta^i w - 2)\right) = 1, \ i = 0, 1, 2 \right\}.$$

Let

$$\widehat{\mathcal{A}}_3 = \left\{ u \ : \ u = \frac{-18}{x^3 - 8},\ (x, y, z, w) \in \mathcal{C}(\mathbb{F}_q),\ yzw \neq 0 \right\}.$$

Clearly, $\widehat{\mathcal{A}}_3 \subset \widetilde{\mathcal{A}}_3$. Now, let $P = (x, y, z, w)$ be an affine point of $\mathcal{C}(\mathbb{F}_q)$ with $yzw \neq 0$. So, $x \neq 0, -\zeta^i, 2\zeta^i$, for $i = 0, 1, 2$. Let $u = \frac{-18}{x^3-8}$. Then, $u \neq 0, 2, 9/4$. So, $u \in \widetilde{\mathcal{A}}_3$ if $2u^2 - 6u + 3 \neq 0$. Thus,

$$\widetilde{\mathcal{A}}_3 = \widehat{\mathcal{A}}_3 \backslash \mathcal{B}_2.$$

In addition, if $2u^2 - 6u + 3 = 0$ then 3 is a quadratic residue in $\mathbb{F}_q$ where $q \equiv 1 \pmod{12}$. And, we have $x^2 + 2x - 2 = 0$. Then, $3x(x+1)(x-2) = 9x^2\beta^2$ and for i=1,2,

$$3\zeta^i x(\zeta^i x + 1)(\zeta^i x - 2) = 9x^2 \zeta^{2i} \beta^{2i-1},$$

where $\beta = (x+1)(2\zeta+1)/3$. Notice that $\beta^2 = -1$. Then

$$\chi_2\left(3\zeta^i x(\zeta^i x + 1)(\zeta^i x - 2)\right) = \begin{cases} 1, & \text{if } i = 0, \\ \chi_2(\beta), & \text{if } i = 1, 2. \end{cases}$$

We note that $\chi_2(\beta) = 1$ if and only if $q \equiv 1 \pmod 8$. Therefore, we have $2u^2 - 6u + 3 = 0$ if and only if $q \equiv 1 \pmod{24}$ and in this case $u \in \widehat{\mathcal{A}}_3$ and $u \notin \widetilde{\mathcal{A}}_3$. So, we have

$$\#\widetilde{\mathcal{A}}_3 = \begin{cases} \#\widehat{\mathcal{A}}_3 - 2, & \text{if } q \equiv 1 \pmod{24} \\ \#\widehat{\mathcal{A}}_3, & \text{if } q \equiv 7, 13 \pmod{24} \\ 0, & \text{if } q \equiv 2 \pmod 3 \end{cases}$$

We consider the map $\tau : \mathcal{C}(\mathbb{F}_q) \to \mathbb{F}_q$ by $\tau(P) = x^3$ that is a $24:1$map for affine points $P = (x, y, z, w)$ with $yzw \neq 0$. And, there are 25 points $P = (x, y, z, w)$ on $\mathcal{C}(\mathbb{F}_q)$ with $yzw = 0$. So, $\#\widehat{\mathcal{A}}_3 = (\#\mathcal{C}(\mathbb{F}_q) - 25)/24$.

In summary, the cardinalities of sets $\widetilde{\mathcal{A}}_i$, for $i = 1, 2, 3$, and $\widetilde{\mathcal{B}}_i$, for $i = 1, 2$, are provided in Table 2.

	$\#\widetilde{\mathcal{A}}_1$	$\#\widetilde{\mathcal{A}}_2$	$\#\widetilde{\mathcal{A}}_3$	$\#\widetilde{\mathcal{B}}_1$	$\#\widetilde{\mathcal{B}}_2$
$q \equiv 1 \pmod{24}$	$\frac{2(q-1)}{3}$	0	$\frac{\#\mathcal{C}(\mathbb{F}_q)-25}{24} - 2$	1	2
$q \equiv 13 \pmod{24}$	$\frac{2(q-1)}{3}$	0	$\frac{\#\mathcal{C}(\mathbb{F}_q)-25}{24}$	1	0
$q \equiv 7 \pmod{12}$	$\frac{2(q-1)}{3}$	0	$\frac{\#\mathcal{C}(\mathbb{F}_q)-25}{24}$	1	0
$q \equiv 2 \pmod 3$	0	$\frac{\#\mathcal{L}(\mathbb{F}_q)-4}{2}$	0	1	0

Table 2: Cardinalities of the sets $\widetilde{\mathcal{A}}_1, \widetilde{\mathcal{A}}_2, \widetilde{\mathcal{A}}_3, \widetilde{\mathcal{B}}_1, \widetilde{\mathcal{B}}_2$.

Now, recall that $I_{\mathrm{T}}(q) = 2J_{\mathrm{T}}(q) - \#\mathcal{T}$. Next, using Theorem 1 and (9), we compute $I_{\mathrm{T}}(q)$ and conclude the proof.

□

4 Doubling Doche-Icart-Kohel curves

4.1 Preliminaries

We consider the curves $\mathbf{E}_{\mathrm{D},u}$ given by (3) over a finite field $\mathbb{F}_q$ of characteristic $p \geq 3$. Notice, $u \notin \{0, 64\}$, since the curve $\mathbf{E}_{\mathrm{D},u}$ is nonsingular. The j-invariant $j(\mathbf{E}_{\mathrm{D},u}) = F(u)$ where

$$F(U) = \frac{(U-48)^3}{U-64}. \tag{10}$$

Let G be the bivariate function

$$\begin{aligned} G(U,V) &= \frac{F(U)-F(V)}{U-V} \\ &= \frac{U^2V + UV^2 - 64U^2 - 208UV - 64V^2 + 9216U + 9216V - 331776}{(U-64)(V-64)}, \end{aligned}$$

and g be the numerator of G given by

$$g(U,V) = U^2V + UV^2 - 64U^2 - 208UV - 64V^2 + 9216U + 9216V - 331776.$$

For $u \in \mathbb{F}_q \backslash \{0, 64\}$ we define the polynomial g_u by $g_u(V) = g(u, V)$, and denote it's discriminant by Δ_u and the set of its $\mathbb{F}_q$ roots by $\mathcal{Z}_u$. So, we have

$$g_u(V) = (u-64)V^2 + (u^2 - 208u + 9216)V + (-64u^2 + 9216u - 331776), \tag{11}$$

$$\Delta_u = u(u-64)(u-48)^2,$$

$$\mathcal{Z}_u = \{v \in \mathbb{F}_q : g_u(v) = 0\}.$$

Recall that two elliptic curves $\mathbf{E}_{\mathrm{D},u}$ and $\mathbf{E}_{\mathrm{D},v}$ are isomorphic over $\overline{\mathbb{F}}_q$ if and only if they have the same j-invariant. In other words, they are $\mathbb{F}_q$-isomorphic if and only if $g_u(v) = g(u, v) = 0$, where $u \neq v$.

For $u \in \mathbb{F}_q \backslash \{0, 64\}$, let

$$\mathcal{J}_u = \left\{ \mathbf{E}_{\mathrm{D},v} : \ v \in \mathbb{F}_q \backslash \{0, 64\}, \ \mathbf{E}_{\mathrm{D},v} \cong_{\overline{\mathbb{F}}_q} \mathbf{E}_{\mathrm{D},u} \right\},$$

$$\mathcal{I}_u = \left\{ \mathbf{E}_{\mathrm{D},v} : \ v \in \mathbb{F}_q \backslash \{0, 64\}, \ \mathbf{E}_{\mathrm{D},v} \cong_{\mathbb{F}_q} \mathbf{E}_{\mathrm{D},u} \right\},$$

and let $\mathcal{N}_q$ be the set of all $\mathbb{F}_q$-isomorphisms between distinct elliptic curves in the family (3) and let $n_q = \#\mathcal{N}_q$. Similarly, let $\overline{n}_q = \#\overline{\mathcal{N}}_q$, where $\overline{\mathcal{N}}_q$ is the set of all $\overline{\mathbb{F}}_q$-isomorphisms between distinct elliptic curves in this family. For $i = 1, 2, 3$, let

$$c_i = \#\left\{ \mathcal{J}_u : u \in \mathbb{F}_q \backslash \{0, 64\}, \ \#\mathcal{J}_u = i \right\}.$$

4.2 Number of $\overline{\mathbb{F}}_q$-isomorphism classes in family (3)

In this section, we compute the number of distinct doubling Doche-Icart-Kohel curves up to twists.

For $u = 0, 64, 48$, where Δ_u equals zero, we have

$$g_0(V) = -64(V-72)^2, \;\; g_{64}(V) = -4096,$$

$$g_{48}(V) = -16(V-48)^2, \;\; \mathcal{J}_{48} = \{\mathbf{E}_{\mathrm{D},48}\}.$$

Also for $u = 72$ (appeared in g_0), we have

$$g_{72}(V) = 8(V-72)V, \;\; \mathcal{J}_{72} = \{\mathbf{E}_{\mathrm{D},72}\}.$$

For all other values of u, $\#\mathcal{J}_u = 3$ if and only if g_u has roots in $\mathbb{F}_q$ that is equivalent to where Δ_u, the discriminant of g_u, is a quadratic residue in $\mathbb{F}_q$.

Lemma 4. *The number c_3 of $\mathbb{F}_q$-isomorphism classes of cardinality three is given by*

$$c_3 = \begin{cases} \dfrac{q-3}{6} & \text{if } q \equiv 0 \pmod 3, \\ \dfrac{q-7}{6} & \text{if } q \equiv 1 \pmod 3, \\ \dfrac{q-5}{6} & \text{if } q \equiv 2 \pmod 3. \end{cases}$$

Proof. In order to find the number of $\overline{\mathbb{F}}_q$-isomorphism classes of cardinality three, we need to enumerate $u \in \mathbb{F}_q \backslash \{0, 64\}$ so that Δ_u is a quadratic residue in $\mathbb{F}_q$. We instead count the number of points (u, z) of the curve

$$u(u-64) = z^2$$

except for the points with $u = 0, 64, 48, 72$, i.e. $\mathcal{O} \cap \mathbb{F}_q^2$, where

$$\mathcal{O} = \{(0,0), (64,0), (72, \pm 24), (48, 16\sqrt{-3})\}.$$

The above curve has $q-1$ points. However, for every $u \in \mathbb{F}_q \backslash \{0, 48, 64, 72\}$ that Δ_u is a quadratic residue, there are two possible values of z. Additionally, every $\overline{\mathbb{F}}_q$-isomorphism class of cardinality three is counted three times. Hence

$$c_3 = \frac{q - 1 - \#(\mathcal{O} \cap \mathbb{F}_q^2)}{6}.$$

□

Theorem 5. *For any finite field $\mathbb{F}_q$ of characteristic $p \geq 3$, for the number $J_{\mathrm{D}}(q)$ of distinct values of the j-invariant of family* (3)*, we have*

$$J_{\mathrm{D}}(q) = \begin{cases} \dfrac{2q-3}{3} & \text{if } q \equiv 0 \pmod 3, \\ \dfrac{2q+1}{3} & \text{if } q \equiv 1 \pmod 3, \\ \dfrac{2q-1}{3} & \text{if } q \equiv 2 \pmod 3. \end{cases}$$

Proof. Since there are only classes of cardinality three and one, we have

$$c_1 = q - 2 - 3c_3.$$

Replacing the values of c_3 according to lemma (4) in the following equation completes the proof.

$$J_{\mathrm{D}}(q) = c_1 + c_3 = q - 2 - 2c_3$$

□

4.3 Number of $\mathbb{F}_q$-isomorphism classes in family (3)

Here, we provide the explicit formulas for the number of $\mathbb{F}_q$-isomorphism classes of doubling Doche-Icart-Kohel curves. The estimate for the number $I_{\mathrm{D}}(q)$ of $\mathbb{F}_q$-isomorphism classes of this family is given by

$$I_{\mathrm{D}}(q) = (19/24)q + O(\sqrt{q}).$$

In order to have $I_{\mathrm{D}}(q)$ explicitly, we compute the number of those $\overline{\mathbb{F}}_q$-isomorphism classes where split into two $\mathbb{F}_q$-isomorphism classes.

Lemma 6. *Let $p \geq 3$. For every $u, v \in \mathbb{F}_q \backslash \{0, 64\}$ that $u \neq v$ and $\mathbf{E}_{\mathrm{D},u} \cong_{\overline{\mathbb{F}}_q} \mathbf{E}_{\mathrm{D},v}$, we have $\mathbf{E}_{\mathrm{D},u} \cong_{\mathbb{F}_q} \mathbf{E}_{\mathrm{D},v}$ if and only if there are elements $a, b \in \mathbb{F}_q$ satisfying*

$$\mathcal{S} \ : \ 32a^2 = \frac{b(b+32)}{b+24}, \tag{12}$$

with

$$\begin{cases} a^2 v = u + 3b, \\ u = \dfrac{-b^2}{b+16}. \end{cases} \tag{13}$$

In fact (u, v) uniquely determines (a^2, b) and vice versa.

Proof. Suppose $\mathbf{E}_{\mathrm{D},u}$ and $\mathbf{E}_{\mathrm{D},v}$ are $\overline{\mathbb{F}}_q$-isomorphic, according to [7] they are isomorphic over $\mathbb{F}_q$, if and only if there is an elliptic curve isomorphism ψ and elements $a, b \in \mathbb{F}_q$ such that

$$\begin{cases} \psi : E_1 \to E_2, \\ \psi(x, y) = (a^2 x + b, a^3 y), \end{cases}$$

and

$$\begin{cases} a^2v = u + 3b, \\ 16a^4v = 16u + 2bu + 3b^2, \\ b(16u + bu + b^2) = 0. \end{cases} \tag{14}$$

Since restrictions on u and v, and properties of ψ imply that $a \neq 0$ and $b \neq 0, -16, -24, -32$, the proof is complete. □

Now, one can see a one to one correspondence between points of $\mathcal{S}$ and $\mathbb{F}_q$-isomorphisms of family (3) except for a few points. The exceptions are the points $(0,0), (0,-32), (\pm i, -16)$ mentioned in the proof of Lemma (6), and $(\pm i, -48), (\pm(-3 \pm i\sqrt{3}) + 2, 8(-3 \pm i\sqrt{3}))$ that are corresponded to isomorphisms of the classes $\mathcal{J}_{72}$ and $\mathcal{J}_{48}$ respectively.

For $p > 3$, the map

$$f : (a, b) \longrightarrow \left(\frac{a(b+24)}{32}, \frac{b}{32} \right)$$

injectively maps $\mathcal{S}$ to

$$\gamma : a'^2 = b'(b' + 1)(b' + 3/4).$$

Furthermore, all the points on γ are images of points on $\mathcal{S}$ except for $(a', b') = (0, -3/4)$ that is corresponded to $b = -24$. Hence we have the set $\mathcal{E}$ of exceptional points on γ

$$\mathcal{E} = f(\{(0,0), (0,-32), (\pm i, -16), (\pm i, -48), (\pm(-3 \pm i\sqrt{3}) + 2, 8(-3 \pm i\sqrt{3}))\}) \cup \{(0, -3/4)\}.$$

If $p = 3$, simplifying $\mathcal{S}$ we have the curve

$$\gamma : a^2 = -b + 1$$

which has q points. But similar to the previous situation, points listed below are not allowed.

$$\mathcal{E} = \{(0,1), (\pm 1, 0), (\pm i, -1)\}$$

In either case

$$n_q = \#\gamma(\mathbb{F}_q) - \#\left(\mathcal{E} \cap \mathbb{F}_q^2\right).$$

Therefore,

$$n_q = \begin{cases} \#\gamma(\mathbb{F}_q) - 3 - 2\chi_2(-1) & \text{if } p = 3, \\ \#\gamma(\mathbb{F}_q) - 3 - 4\chi_2(-1) - 4\chi_2(-3) & \text{if } p \neq 3. \end{cases}$$

So,

$$n_q = \begin{cases} q - 5 & \text{if } q = 3^{2k}, \\ q - 3 & \text{if } q = 3^{2k+1}, \\ \#\gamma(\mathbb{F}_q) - 11 & \text{if } q \equiv 1 \pmod{12}, \\ \#\gamma(\mathbb{F}_q) - 7 & \text{if } q \equiv 5, 7 \pmod{12}, \\ \#\gamma(\mathbb{F}_q) - 3 & \text{if } q \equiv 11 \pmod{12}. \end{cases}$$

In lemma (6), whether $\mathbf{E}_{\mathrm{D},u}$ and $\mathbf{E}_{\mathrm{D},v}$ are $\mathbb{F}_q$-isomorphic or not, equations (12) and (13) have two pairs of solutions $(\pm a, b) \in \overline{\mathbb{F}}_q^2$ that are correspondent to the two $\overline{\mathbb{F}}_q$-isomorphisms from $\mathbf{E}_{\mathrm{D},u}$ to $\mathbf{E}_{\mathrm{D},v}$. Therefore, there are twelve $\overline{\mathbb{F}}_q$-isomorphisms between distinct curves in a cardinality three $\overline{\mathbb{F}}_q$-isomorphism class (two isomorphisms for each two-permutation of three $\overline{\mathbb{F}}_q$-isomorphic curve). So

$$\overline{n}_q = 12c_3.$$

Since every two $\overline{\mathbb{F}}_q$-isomorphic curve are either $\mathbb{F}_q$-isomorphic or nontrivial quadratic twists of each other, every $\overline{\mathbb{F}}_q$-isomorphism class of size three either splits into one or two $\mathbb{F}_q$-isomorphism classes. Therefore, there are either four or twelve $\mathbb{F}_q$-isomorphisms between distinct curves in every $\overline{\mathbb{F}}_q$-isomorphism class of size three. In other words, every eight $\overline{\mathbb{F}}_q$-isomorphisms that are not $\mathbb{F}_q$-isomorphisms, increase the number of $\mathbb{F}_q$-isomorphism classes by one. Hence

$$I_{\mathrm{D}}(q) - J_{\mathrm{D}}(q) = \frac{\overline{n}_q - n_q}{8}. \tag{15}$$

Theorem 7. *For any finite field $\mathbb{F}_q$ of characteristic $p \geq 3$, for the number $I_{\mathrm{D}}(q)$ of $\mathbb{F}_q$-isomorphism classes of the family* (3)*, we have*

$$I_{\mathrm{D}}(q) = \begin{cases} (11q+1)/12 - N/8 & \text{if } q \equiv 1 \pmod{12}, \\ (11q-7)/12 - N/8 & \text{if } q \equiv 5 \pmod{12}, \\ (11q-5)/12 - N/8 & \text{if } q \equiv 7 \pmod{12}, \\ (11q-13)/12 - N/8 & \text{if } q \equiv 11 \pmod{12}, \\ (19q-27)/24 & \text{if } q = 3^{2k}, \\ (19q-33)/24 & \text{if } q = 3^{2k+1}, \end{cases}$$

where N is the number of $\mathbb{F}_q$-rational points on the elliptic curve

$$L : Y^2 = X(X+1)(X+3/4).$$

Proof. According to (15),

$$I_{\mathrm{D}}(q) = J_{\mathrm{D}}(q) + \frac{\overline{n}_q - n_q}{8}.$$

Replacing the values of n_q and $\overline{n}_q$ completes the proof. □

5 Conclusion

In this paper, we give explicit formulas for the number of distinct elliptic curves over a finite field, up to isomorphism, in two families (2) and (3) of curves introduced by Doche, Icart and Kohel. For any finite field $\mathbb{F}_q$ of characteristic $p \geq 5$, for the number $J_{\mathrm{T}}(q)$ of $\overline{\mathbb{F}}_q$-isomorphism classes of the family (2), we have [5, Theorem 9]

$$J_{\mathrm{T}}(q) = \begin{cases} \left\lfloor \dfrac{3q+1}{4} \right\rfloor & \text{if } q \equiv 1 \pmod{3}, \\ \dfrac{q-1}{2} & \text{if } q \equiv 2 \pmod{3}. \end{cases}$$

We recall [2] that for $\mathbb{F}_q$ of characteristic $p \geq 5$ the family (2) is the family of elliptic curves defined over $\mathbb{F}_q$ with a rational 3-torsion subgroup (up to twists). In other words the formulas provide the number of $\overline{\mathbb{F}}_q$-isomorphism classes of the family of elliptic curves over $\mathbb{F}_q$ with a rational 3-torsion point. In addition, an elliptic curve with a point of order 3 is birationally equivalent to a generalized Hessian curve. The exact number of distinct elliptic curves with a point of order 3 up to isomorphism is also provided by [3]. And, the the number of $\mathbb{F}_q$-isomorphism classes of the family of elliptic curves over $\mathbb{F}_q$ with a rational 3-torsion point is given by

$$\begin{cases} \lfloor (3(q+3)/4 \rfloor & \text{if } q \equiv 1 \pmod 3, \\ q-1 & \text{if } q \equiv 2 \pmod 3. \end{cases}$$

The interesting result of this paper is Theorem (3) providing the estimate for the number $I_{\mathrm{T}}(q)$ of $\mathbb{F}_q$-isomorphism classes of this family by

$$I_{\mathrm{T}}(q) = \begin{cases} (79/96)q + O(\sqrt{q}) & \text{if } q \equiv 1 \pmod 3, \\ (3/4)q + O(\sqrt{q}) & \text{if } q \equiv 2 \pmod 3. \end{cases}$$

Therefore, the family of Doche, Icart and Kohel does not cover all $\mathbb{F}_q$-isomorphism classes of elliptic curves with a point of order 3. And, the more intrusting point is that the number $I_{\mathrm{T}}(q)$ is estimated by Hasse-Weil bound.

For the family of doubling Doche, Icart and Kohel curves over any finite field $\mathbb{F}_q$ of characteristic $p \geq 5$, Theorem (5) provides the number $J_{\mathrm{D}}(q)$ of $\overline{\mathbb{F}}_q$-isomorphism classes of this family (3) by

$$J_{\mathrm{D}}(q) = \left\lfloor \frac{2q+1}{3} \right\rfloor.$$

This family represents elliptic curves defined over $\mathbb{F}_q$ with a rational 2-torsion subgroup (up to twists). Moreover, the number of $\mathbb{F}_q$-isomorphism classes of the family of elliptic curves over $\mathbb{F}_q$ with a rational 2-torsion point is given by

$$\begin{cases} 2\left\lfloor \frac{2q+1}{3} \right\rfloor + 2 & \text{if } q \equiv 1 \pmod 4, \\ 2\left\lfloor \frac{2q+1}{3} \right\rfloor & \text{if } q \not\equiv 1 \pmod 4. \end{cases}$$

And, Theorem (3) gives the estimate for the number $I_{\mathrm{D}}(q)$ of $\mathbb{F}_q$-isomorphism classes of this family by

$$I_{\mathrm{D}}(q) = (19/24)q + O(\sqrt{q}).$$

In conclusion, we studied interesting examples of families of elliptic curves with a small subgroup where the number of $\mathbb{F}_q$-isomorphism classes of the family is not exactly given by the size of the finite field and it is estimated by Hasse-Weil bound.

References

[1] R. Avanzi, H. Cohen, C. Doche, G. Frey, T. Lange, K. Nguyen and F. Vercauteren, *Handbook of elliptic and hyperelliptic curve cryptography*, CRC Press, 2005.

[2] C. Doche, T. Icart and D. R. Kohel, 'Efficient scalar multiplication by isogeny decompositions', *PKC'2006*, LNCS, vol. 3958, Springer-Verlag, 2006, 191–206.

[3] R. R. Farashahi, 'On the Number of Distinct Legendre, Jacobi, Hessian and Edwards Curves (Extended Abstract)', Workshop on Coding theory and Cryptology—WCC 2011, (2011), 37–46.

[4] R. R. Farashahi, D. Moody and Hongfeng WU, 'Isomorphism classes of Edwards and twisted Edwards curves over finite fields', *Finite Fields and Their Applications*, **18**(3), (2012), 597–612.

[5] R. R. Farashahi and I. E. Shparlinski, 'On the number of distinct elliptic curves in some families', *Designs, Codes and Cryptography*, **54**(1), (2010), 83–99.

[6] E. Howe, 'On the group orders of elliptic curves over finite fields', *Compositio Mathematica*, **85**(2), (1993), 229–247.

[7] J. H. Silverman, *The arithmetic of elliptic curves*, Springer-Verlag, Berlin, 1995.

[8] L. C. Washington, *Elliptic curves: Number theory and Cryptography*, CRC Press, 2008.